


450nm

• wave length 750nm Shorter wave length images have stronger strange pattern. Longer wave length images have stronger fringes.


750nm




Steps for Measuring Tree Ring Patterns

- Stacking
- Masking
- Polynomial fitting and normalization
- Correcting other patterns
- Determining Tree Ring Center
- Measuring Tree Ring Pattern
- Testing Tree Ring Cancelation
- Calculating Displacement by Tree Ring
- Calculating Spurious Shear by Tree Ring

Masking




Masking




Polynomial fitting and normalization

- 7th order polynomial fitting for each channels and normalize.
- AVE = 1.0
- STD = 0.004





Polynomial fitting and normalization


- Tree Ring can be seen from smoothed image.
- some other patterns which don't look like tree ring are visible.
- grid pattern(pixel size?)




• sloped line pattern(?).











Average counts horizontal




- Lines of the slopes can be fitted by y = 0.15x.
- fitting each lines and normalize


Fig.8-2




• large scale fluctuation subtraction in Fourier space. No subtraction



• large scale fluctuation subtraction in Fourier space. Subtraction $|kx| \le 1$ and $|ky| \le 1$



• large scale fluctuation subtraction in Fourier space. Subtraction $|kx| \le 2$ and $|ky| \le 2$




• large scale fluctuation subtraction in Fourier space. Subtraction $|kx| \le 3$ and $|ky| \le 3$




• large scale fluctuation subtraction in Fourier space. Subtraction $|kx| \le 4$ and $|ky| \le 4$




• large scale fluctuation subtraction in Fourier space. Subtraction $|kx| \le 5$ and $|ky| \le 5$




• large scale fluctuation subtraction in Fourier space. Subtraction $|kx| \le 6$ and $|ky| \le 6$




• large scale fluctuation subtraction in Fourier space. Subtraction $|kx| \le 7$ and $|ky| \le 7$




• large scale fluctuation subtraction in Fourier space. Subtraction $|kx| \le 8$ and $|ky| \le 8$





- Measuring positions of lines of tree ring and getting center of the ring by least square fitting.
- (Temporary) Center [4575, -375]




Center











Testing Tree Ring Cancelation



Before Cancelation (Smoothed image)

After Cancelation (Smoothed image)


Measuring Tree Ring Pattern



Calculating Displacement by Tree Ring

Andres's equation for estimating displacement from tree ring effect for flat.


$$f(r) = \frac{1}{r} \int rw(r)dr$$



Calculating Spurious Shear by Tree Ring


A formula for estimating spurious shear from concentric displacement

$$\kappa = \frac{1}{2} \left(\frac{\partial f(r)}{\partial r} + \frac{f(r)}{r} \right)$$
$$\gamma = \frac{1}{2} \left(\frac{\partial f(r)}{\partial r} - \frac{f(r)}{r} \right)$$




Tree Ring on LSST CCD


- ITL CCD
- wave length 750nm Shorter wave length images have stronger strange pattern. Longer wave length images have stronger fringes.
- Almost same steps as e2cv CCD, except correcting bamboo patterns.




Bamboo pattern



Correcting Bamboo Pattern




Average counts vertical lines




Average counts horizontal lines

Correcting Bamboo Pattern



Tree Ring on LSST CCD



Conclusion

I measured the tree ring pattern on two LSST CCDs. There still are some issues for more precise measurement, determination of the center, removing noise peaks and so on. However, the results are enough to estimate typical scale of the tree ring pattern. The estimated scale is 0.01% for flux fluctuation, 0.001 pixel for displacement and 0.005% for spurious shear from both CCDs. These are 1/50 scale of them of DEC and small enough to be ignored.

